Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Association of Ambient Air Pollution with Depressive and Anxiety Symptoms in Older Adults: Results from the NSHAP Study

Vivian C. Pun, Justin Manjourides, and Helen Suh

Table of Contents

Figure S1 Flowchart of participants through the NSHAP study

Figure S2 ORs (95% CIs) for mental illness per 5 μ g/m³ increment in PM_{2.5} levels over extended range of exposure moving averages in multivariable model.

Table S1 Assessment of depression, anxiety, and perceived stress measures in the NSHAP study

Table S2 Description of covariates in the NSHAP study

Table S3 Interquartile range of PM_{2.5} moving averages

Table S4 Characteristics of NSHAP study participants by waves and PM_{2.5} pollution categories

Table S5 ORs (95% CIs) for mental disorders per 5 μ g/m³ increment in PM_{2.5} levels over preceding 30 days moving average – evaluating potential confounding

Table S6 ORs (95% CI) for mental illness per 5 μ g/m³ increment in PM_{2.5} levels over various moving averages – comparison of imputation methods using multivariable models

Table S7 ORs (95% CIs) for mental illness per 5 μ g/m³ increment in PM_{2.5} levels over various preceding days moving averages – Restricting to only individuals who participated in both waves

Table S8 Percent change (95% CIs) in scores of mental health indicators per 5 μ g/m³ increment in PM_{2.5} levels over preceding days moving averages

Table S9 Sensitivity analyses of the ORs (95% CIs) for mental disorders per 5 μ g/m³ increment in PM_{2.5} levels over various preceding days moving averages in multivariable models

Table S10 Additional effect modification analysis of the association of mental illness with 5 $\mu g/m^3$ increment in PM_{2.5} levels over preceding 30 days moving average in multivariable models with interaction terms for the potential modifier